Neuroscience Letters, 99 (1989) 356–359
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abad, F., Feria, M. and Boada, J., Chronic amitriptyline decreases autotomy following dorsal rhizotomy in rats, 187
Agmon, A. and Connors, B.W., Repetitive burst-firing neurons in the deep layers of mouse somatosensory cortex, 137
Agnati, L.F., see Tinner, B., 44
Akiguchi, I., see Shigematsu, K., 18
Alzheimer, C., Sutor, B. and Ten Bruggencate, G., Transient and selective blockade of adenosine A1-receptors by 8-cyclopentyl-1,3-dipropylxanthine (DPCPX) causes sustained epileptiform activity in hippocampal CA3 neurons of guinea pigs, 107
Ammermüller, J., see Schlemermeyer, E., 73
Andersson, K., see Tinner, B., 44
Anthony, B.L., Dennison, R.L. and Aronstam, R.S., Disruption of muscarinic receptor—G protein coupling is a general property of liquid volatile anesthetics, 191
Arimatsu, Y. and Miyamoto, M., Colocalization of NGF receptor immunoreactivity and [^{3}H]GABA uptake activity in developing rat septum/diagonal band neurons in vitro, 39
Aronstam, R.S., see Anthony, B.L., 191

Barakat, I. and Droz, B., Maintenance of neuronal expression of calbindin by a muscular extract in cultures of chick dorsal root gangion cells, 1
Barber, A., Harting, J. and Wolf, H.-P., Antinociceptive effects of the 5-HT$_2$ antagonist ritanserin in rats: evidence for an activation of descending monoamingergic pathways in the spinal cord, 234
Basbaum, A.I., see Weill-Fugazza, J., 181
Batini, C., Buisseret-Delmas, C., Coimpoint, C. and Daniel, H., The GABAergic neurones of the cerebellar nuclei in the rat: projections to the cerebellar cortex, 251
Beattie, G.M., see Lakshmanan, J., 263
Beck, S.G., 5-Carboxyamidotryptamine mimics only the 5-HT elicited hyperpolarization of hippocampal pyramidal cells via 5-HT$_1$ receptor, 101
Ben-Ari, Y., see Represa, A., 345
Ben-Ari, Y., see Represa, A., 351
Ben-Ari, Y., see Represa, A., 61
Bettaney, J., see Dray, A., 50
Boada, J., see Abad, F., 187
Bockaert, J., see Rassendren, F.-A., 333
Boegman, R.J. and Metcalf, R.H., Effect of unilateral decortication on acetylcholine release from the nucleus basalis, 219
Böhmer, G., see Schmid, K., 305
Brewer, G.J. and Cotman, C.W., NMDA receptor regulation of neuronal morphology in cultured hippocampal neurons, 268
Buisseret-Delmas, C., see Batini, C., 251
Burns, C., see Lakshmanan, J., 263

Chamak, B., see Mallat, M., 12
Chen, K.Y., see Hsu, L., 257
Chevallier, J., see Grandier-Vazeille, X., 30
Chizhmakov, I.V., Kiskin, N.I., Krishtal, O.A. and Tsyndrenko, A.Ya., Glycine action on *N*-methyl-D-aspartate receptors in rat hippocampal neurons, 131
Clark, G.D., see Zorumski, C.F., 293
Clifford, D.B., see Zorumski, C.F., 293
Coffey, A.R. and Kevetter, G.A., Ipsilateral and contralateral projections to the trochlear nucleus arise from different subdivisions in the vestibular nuclei, 274
Coimpoint, C., see Batini, C., 251
Connors, B.W., see Agmon, A., 137
Cotman, C.W., see Brewer, G.J., 268

Daniel, H., see Batini, C., 251
De Vries, J.B., see Westerink, B.H.C., 197
De la Garza, R., Freedman, R. and Hoffer, B.J., κ-Bungarotoxin blockade of nicotine electrophysiological actions in cerebellar Purkinje neurons, 95
Deguchi, M., Niwa, M., Shigematsu, K., Fujii, T., Namba, K. and Ozaki, M., Specific [^{125}I]Bolton–Hunter substance P binding sites in human rat skin, 287
Demeulemeester, H., Vandesande, F., Orban, G.A., Heizmann, C.W. and Pochet, R., Calbinding D-28K and parvalbumin immunoreactivity is confined to two separate neuronal subpopulations in the cat visual cortex, whereas partial coexistence is shown in the dorsal lateral geniculate nucleus, 6

0304-3940/89/$ 03.50 © 1989 Elsevier Scientific Publishers Ireland Ltd.

Neuroscience Letters, 99 (1989) 356–359
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abad, F., Feria, M. and Boada, J., Chronic amitriptyline decreases autotomy following dorsal rhizotomy in rats, 187
Agmon, A. and Connors, B.W., Repetitive burst-firing neurons in the deep layers of mouse somatosensory cortex, 137
Agnati, L.F., see Tinner, B., 44
Akiguchi, I., see Shigematsu, K., 18
Alzheimer, C., Sutor, B. and Ten Bruggencate, G., Transient and selective blockade of adenosine A1-receptors by 8-cyclopentyl-1,3-dipropylxanthine (DPCPX) causes sustained epileptiform activity in hippocampal CA3 neurons of guinea pigs, 107
Ammermüller, J., see Schlemermeyer, E., 73
Andersson, K., see Tinner, B., 44
Anthony, B.L., Dennison, R.L. and Aronstam, R.S., Disruption of muscarinic receptor—G protein coupling is a general property of liquid volatile anesthetics, 191
Arimatsu, Y. and Miyamoto, M., Colocalization of NGF receptor immunoreactivity and [^{3}H]GABA uptake activity in developing rat septum/diagonal band neurons in vitro, 39
Aronstam, R.S., see Anthony, B.L., 191

Barakat, I. and Droz, B., Maintenance of neuronal expression of calbindin by a muscular extract in cultures of chick dorsal root gangion cells, 1
Barber, A., Harting, J. and Wolf, H.-P., Antinociceptive effects of the 5-HT$_2$ antagonist ritanserin in rats: evidence for an activation of descending monoamingergic pathways in the spinal cord, 234
Basbaum, A.I., see Weill-Fugazza, J., 181
Batini, C., Buisseret-Delmas, C., Coimpoint, C. and Daniel, H., The GABAergic neurones of the cerebellar nuclei in the rat: projections to the cerebellar cortex, 251
Beattie, G.M., see Lakshmanan, J., 263
Beck, S.G., 5-Carboxyamidotryptamine mimics only the 5-HT elicited hyperpolarization of hippocampal pyramidal cells via 5-HT$_1$ receptor, 101
Ben-Ari, Y., see Represa, A., 345
Ben-Ari, Y., see Represa, A., 351
Ben-Ari, Y., see Represa, A., 61
Bettaney, J., see Dray, A., 50
Boada, J., see Abad, F., 187
Bockaert, J., see Rassendren, F.-A., 333
Boegman, R.J. and Metcalf, R.H., Effect of unilateral decortication on acetylcholine release from the nucleus basalis, 219
Böhmer, G., see Schmid, K., 305
Brewer, G.J. and Cotman, C.W., NMDA receptor regulation of neuronal morphology in cultured hippocampal neurons, 268
Buisseret-Delmas, C., see Batini, C., 251
Burns, C., see Lakshmanan, J., 263

Chamak, B., see Mallat, M., 12
Chen, K.Y., see Hsu, L., 257
Chevallier, J., see Grandier-Vazeille, X., 30
Chizhmakov, I.V., Kiskin, N.I., Krishtal, O.A. and Tsyndrenko, A.Ya., Glycine action on *N*-methyl-D-aspartate receptors in rat hippocampal neurons, 131
Clark, G.D., see Zorumski, C.F., 293
Clifford, D.B., see Zorumski, C.F., 293
Coffey, A.R. and Kevetter, G.A., Ipsilateral and contralateral projections to the trochlear nucleus arise from different subdivisions in the vestibular nuclei, 274
Coimpoint, C., see Batini, C., 251
Connors, B.W., see Agmon, A., 137
Cotman, C.W., see Brewer, G.J., 268

Daniel, H., see Batini, C., 251
De Vries, J.B., see Westerink, B.H.C., 197
De la Garza, R., Freedman, R. and Hoffer, B.J., κ-Bungarotoxin blockade of nicotine electrophysiological actions in cerebellar Purkinje neurons, 95
Deguchi, M., Niwa, M., Shigematsu, K., Fujii, T., Namba, K. and Ozaki, M., Specific [^{125}I]Bolton–Hunter substance P binding sites in human rat skin, 287
Demeulemeester, H., Vandesande, F., Orban, G.A., Heizmann, C.W. and Pochet, R., Calbinding D-28K and parvalbumin immunoreactivity is confined to two separate neuronal subpopulations in the cat visual cortex, whereas partial coexistence is shown in the dorsal lateral geniculate nucleus, 6

0304-3940/89/$ 03.50 © 1989 Elsevier Scientific Publishers Ireland Ltd.

Dennison, R.L., see Anthony, B.L., 191
Dichgans, J., see Mai, N., 340
Diener, H.-C., see Mai, N., 340
Dietzel, I.D., see Gottmann, K., 90
Dolphin, A.C., see McGuirk, S.M., 85
Dray, A., Bettaney, J. and Forster, P., Capsaicin desensitization of peripheral nociceptive fibres does not impair sensitivity to other noxious stimuli, 50
Dreyer, D., Lagrange, A., Grothe, C. and Unsicker, K., Basic fibroblast growth factor prevents ontogenetic neuron death in vivo, 35
Droz, B., see Barakat, I., 1
Duvert, M., see Grandier-Vazeille, X., 30

Errchidi, S., see Monteau, R., 311

Feria, M., see Abad, F., 187
Fisher, D.A., see Lakshmanan, J., 263
Fonnum, F., see Fykse, E.M., 300
Forster, P., see Dray, A., 50
Freedman, R., see De la Garza, R., 95
Fujii, T., see Deguchi, M., 287
Fuxe, K., see Tinner, B., 44
Fykse, E.M. and Fonnum, F., Regional distribution of γ-aminobutyrate and L-glutamate uptake into synaptic vesicles isolated from rat brain, 300

Garthwaite, G. and Garthwaite, J., Quisqualate neurotoxicity: a delayed, CNQX-sensitive process triggered by a CNQX-insensitive mechanism in young rat hippocampal slices, 113
Garthwaite, J., see Garthwaite, G., 113
Gauthier, P., see Monteau, R., 311
Gebauer, K., see Schmid, K., 305
Godefroy, F., see Weill-Fugazza, J., 181
Gottmann, K., Dietzel, I.D., Lux, H.D. and Ruedel, C., Proton-induced Na^+ current develops prior to voltage-dependent Na^+ and Ca^{2+} currents in neuronal precursor cells from chick dorsal root ganglion, 90
Grandier-Vazeille, X., Duvert, M. and Chevallier, J., The use of auto-anti-idiotypes for the visualization of acetylcholine receptors in an invertebrate skeletal muscle, 30
Grothe, C., see Dreyer, D., 35
Gustow, E., see Ved, H.S., 203

Hämäläinen, M., see Hari, R., 164
Han, Z., see Ju, G., 246
Hanada, M., see Tamiya, R., 241
Hannah, R.S., Hayden, L.J. and Roth, S.H., Hydrogen sulfide exposure alters the amino acid content in developing rat CNS, 323
Harding, R., see Rees, S., 157
Hari, R., Joutsiniemi, S.L., Hämäläinen, M. and Vilkman, V., Neuromagnetic responses of human auditory cortex to interruptions in a steady rhythm, 164
Harting, J., see Barber, A., 234
Hashizume, K., see Kanda, K., 153
Hayden, L.J., see Hannah, R.S., 323
Hayek, A., see Lakshmanan, J., 263
Heizmann, C.W., see Demeulemeester, H., 6
Herranz, A.S., see Herreras, O., 119
Herreras, O., Menéndez, A., Herranz, A.S., Solis, J.M. and Martín del Río, R., Synaptic transmission at the Schaffer-CA1 synapse is blocked by 6,7-dinitroquinoxaline-2,3-dione. An in vivo brain dialysis study in the rat, 119
Hersh, L., see Tinner, B., 44
Hilaire, G., see Monteau, R., 311
Hilliges, M., see Johansson, O., 281
Hoffer, B.J., see De la Garza, R., 95
Hsu, L., Jeng, A.Y. and Chen, K.Y., Induction of neurite outgrowth from chick embryonic ganglia explants by activators of protein kinase C, 257
Hunt, A., see Patel, A.J., 223

Ichinose, H., see Naoi, M., 317
Imura, H., see Koshiyama, H., 328
Jansson, A., see Tinner, B., 44
Jeng, A.Y., see Hsu, L., 257
Johansson, O., Hilliges, M. and Stahle-Bäckdahl, M., Intraepidermal neuron-specific enolase (NSE)-immunoreactive nerve fibres: evidence for sprouting in uremic patients on maintenance hemodialysis, 281
Johnson, R.F., see Morin, L.P., 55
Joutsiniemi, S.L., see Hari, R., 164
Ju, G. and Han, Z., Coexistence of corticotropin releasing factor and neurotensin within oval nucleus neurons in the bed nuclei of the stria terminalis in the rat, 246

Kameyama, M., see Shigematsu, K., 18
Kamo, H., see Shigematsu, K., 18
Kanda, K., Sato, H., Hashizume, K. and Yamada, J., The effects of blocking nerve conduction on retrograde HRP labeling of rat motoneuron, 153
Kato, Y., see Koshiyama, H., 328
Kawai, Y., see Tamiya, R., 241

Kevetter, G.A., see Coffey, A.R., 274
Kijima, H., see Tanabe, N., 147
Kimura, H., see Shigematsu, K., 18
Kiskin, N.I., see Chizhmakov, I.V., 131
Kitamura, T., Yamada, Y. and Sato, H., Axon collaterals of spinocerebellar fibers terminate in the parabrachial nucleus of the rat, 24
Koerber, H.R., Seymour, A.W. and Mendell, L.M., Mismatches between peripheral receptor type and central projections after peripheral nerve regeneration, 67
Köhler, Ch., see Tinner, B., 44
Kohler, K., see Weiler, R., 214
Kolbinger, W., see Weiler, R., 214
Konopka, L.M. and Parsons, R.L., Characteristics of the galanin-induced depolarization of mudpuppy parasympathetic postganglionic neurons, 142
Koshiyama, H., Kato, Y., Shimatsu, A. and Imura, H., Effects of clonidine and naloxone on the dopamine levels in the rat mediobasal hypothalamus measured by in vivo voltammetry, 328
Krishtal, O.A., see Chizhmakov, I.V., 131

Lagrange, A., see Dreyer, D., 35
Lakshmanan, J., Beattie, G.M., Hayek, A., Burns, C. and Fisher, D.A., Biological actions of 53 kDa nerve growth factor as studied by a blot and culture technique, 263
Lanthorn, T., see Rader, R.K., 125
Le Gal La Salle, G., see Represa, S., 345
Lory, P., see Rassendren, F.-A., 333
Lux, H.D., see Gottmann, K., 90

Mai, N., Diener, H.-C. and Dichgans, J., On the role of feedback in maintaining constant grip force in patients with cerebellar disease, 340
Maksay, G. and Simonyi, M., Correlation between regional and kinetic heterogeneities of β-carboline/benzodiazepine receptor binding in rat brain, 229
Mallat, M. and Chamak, B., Lineage relationship between oligodendrocytes and brain macrophages?, 12
Martín del Río, R., see Herreras, O., 119
McGuirk, S.M., Vallis, Y., Pasternak, C.A. and Dolphin, A.C., Bradykinin enhances excitability in cultured rat sensory neurones by a GTP-dependent mechanism, 85
Mendell, L.M., see Koerber, H.R., 67
Menéndez, A., see Herreras, O., 119
Metcalf, R.H., see Boegman, R.J., 219
Miyamoto, M., see Arimatsu, Y., 39
Monteau, R., Errchidi, S., Gauthier, P., Hilaire, G. and Rega, P., Pneumotaxic centre and apneustic breathing: interspecies differences between rat and cat, 311
Moore, R.Y., see Morin, L.P., 55
Morin, L.P., Johnson, R.F. and Moore, R.Y., Two brain nuclei controlling circadian rhythms are identified by GFAP immunoreactivity in hamsters and rats, 55

Nagatsu, T., see Naoi, M., 317
Namba, K., see Deguchi, M., 287
Naoi, M., Takahashi, T., Ichinose, H., Wakabayashi, K., Sugimura, T. and Nagatsu, T., Uptake of heterocyclic amines, Trp-P-1 and Trp-P-2, into clonal rat pheochromocytoma PC12h cells by dopamine uptake system, 317
Nargeot, J., see Rassendren, F.-A., 333
Narici, L., Romani, G.L., Traversa, R. and Rossini, P.M., Neuromagnetic imaging studies discriminate proprioceptive and cutaneous cortical inputs during median nerve stimulation in man, 169
Narita, N., see Tamiya, R., 241
Niwa, M., see Deguchi, M., 287
Northcutt, R.G. and Plassmann, W., Electrosensory activity in the telencephalon of the axolotl, 79

Orban, G.A., see Demeulemeester, H., 6
Ozaki, M., see Deguchi, M., 287

Parsons, R.L., see Konopka, L.M., 142
Pasternak, C.A., see McGuirk, S.M., 85
Patel, A.J. and Hunt, A., Regulation of production by primary cultures of rat forebrain astrocytes of a trophic factor important for the development of cholinergic neurons, 223
Pieringer, R.A., see Ved, H.S., 203
Pin, J.-P., see Rassendren, F.-A., 333
Plassmann, W., see Northcutt, R.G., 79
Pochet, R., see Demeulemeester, H., 6
Proske, U., see Rees, S., 157

Rader, R.K. and Lanthorn, T., Experimental ischemia induces a persistent depolarization blocked by decreased calcium and NMDA antagonists, 125
Rassendren, F.-A., Lory, P., Pin, J.-P., Bockaert, J. and Nargeot, J., A specific quisqualate agonist inhibits kainate responses induced in *Xenopus* oocytes injected with rat brain RNA, 333

Rees, S., Proske, U. and Harding, R., Conduction velocity and fibre diameter of the peroneal nerve in normal and growth retarded fetal sheep, 157
Rega, P., see Monteau, R., 311
Represa, A., Le Gal La Salle, G. and Ben-Ari, Y., Hippocampal plasticity in the kindling model of epilepsy in rats, 345
Represa, A., Robain, O., Tremblay, E. and Ben-Ari, Y., Hippocampal plasticity in childhood epilepsy, 351
Represa, A., Tremblay, E. and Ben-Ari, Y., Transient increase of NMDA-binding sites in human hippocampus during development, 61
Robain, O., see Represa, A., 351
Romani, G.L., see Narici, L., 169
Rossini, P.M., see Narici, L., 169
Roth, S.H., see Hannah, R.S., 323
Ruedel, C., see Gottmann, K., 90
Russell, R.D., see Stripling, J.S., 208

Sato, H., see Kanda, K., 153
Sato, H., see Kitamura, T., 24
Schlemermeyer, E., Schütte, M. and Ammermüller, J., Immunohistochemical and electrophysiological evidence that locust ocellar photoreceptors contain and release histamine, 73
Schmid, K., Böhmer, G. and Gebauer, K., $GABA_B$ receptor mediated effects on central respiratory system and their antagonism by phaclofen, 305
Schütte, M., see Schlemermeyer, E., 73
Seymour, A.W., see Koerber, H.R., 67
Shefchyck, S.J., The effects of lumbosacral deafferentation on pontine micturition centre-evoked voiding in the decerebrate cat, 175
Shigematsu, K., Kamo, H., Akiguchi, I., Kameyama, M. and Kimura, H., Neovascularization of transplanted central nervous tissue suspensions: an immunohistochemical study with laminin, 18
Shigematsu, K., see Deguchi, M., 287
Shimatsu, A., see Koshiyama, H., 328
Simonyi, M., see Maksay, G., 229
Solis, J.M., see Herreras, O., 119
Ståhle-Bäckdahl, M., see Johansson, O., 281
Stripling, J.S. and Russell, R.D., Twenty-four-hour post-seizure inhibition during limbic kindling requires seizure generalization, 208
Sugimura, T., see Naoi, M., 317
Sutor, B., see Alzheimer, C., 107

Takagi, Y., see Tamiya, R., 241
Takahashi, T., see Naoi, M., 317
Tamiya, R., Hanada, M., Narita, N., Kawai, Y., Tohyama, M. and Takagi, Y., Neuropeptide Y afferents have synaptic interactions with histaminergic (histidine decarboxylase-immunoreactive) neurons in the rat brain, 241
Tanabe, N. and Kijima, H., Both augmentation and potentiation occur independently of internal Ca^{2+} at the frog neuromuscular junction, 147
Ten Bruggencate, G., see Alzheimer, C., 107
Thio, L.L., see Zorumski, C.F., 293
Tinner, B., Fuxe, K., Köhler, Ch., Hersh, L., Andersson, K., Jansson, A. and Agnati, L.F., Evidence for the existence of a population of arcuate neurons costoring choline acetyltransferase and tyrosine hydroxylase immunoreactivities in the male rat, 44
Tohyama, M., see Tamiya, R., 241
Traversa, R., see Narici, L., 169
Tremblay, E., see Represa, A., 351
Tremblay, E., see Represa, A., 61
Tsyndrenko, A.Ya., see Chizhmakov, I.V., 131

Unsicker, K., see Dreyer, D., 35

Vallis, Y., see McGuirk, S.M., 85
Vandesande, F., see Demeulemeester, H., 6
Ved, H.S., Gustow, E. and Pieringer, R.A., Effect of hydrocortisone on myelin basic protein in developing primary brain cultures, 203
Vilkman, V., see Hari, R., 164

Wakabayashi, K., see Naoi, M., 317
Weiler, R., Kolbinger, W. and Kohler, K., Reduced light responsiveness of the cone pathway during prolonged darkness does not result from an increase of dopaminergic activity in the fish retina, 214
Weill-Fugazza, J., Godefroy, F. and Basbaum, A.I., Effect of deafferentation on the levels of uric acid in the spinal cord of the rat, 181
Westerink, B.H.C. and De Vries, J.B., On the mechanism of neuroleptic induced increase in striatal dopamine release: brain dialysis provides direct evidence for mediation by autoreceptors localized on nerve terminals, 197
Wolf, H.-P., see Barber, A., 234

Yamada, J., see Kanda, K., 153
Yamada, Y., see Kitamura, T., 24

Zorumski, C.F., Thio, L.L., Clark, G.D. and Clifford, D.B., Calcium influx through *N*-methyl-D-aspartate channels activates a potassium current in postnatal rat hippocampal neurons, 293